1	Mutability of mononucleotide repeats, not oxidative stress, explains the discrepancy between
2	laboratory-accumulated mutations and the natural allele-frequency spectrum in C. elegans
3	
4	Moein Rajaei ¹ , Ayush Shekhar Saxena ¹ , Lindsay M. Johnson ¹ , Michael C. Snyder ¹ , Timothy A.
5	Crombie ^{1,2} , Robyn E. Tanny ² , Erik C. Andersen ² , Joanna Joyner-Matos ³ , and Charles F. Baer ^{1,4}
6	
7	1- Department of Biology, University of Florida, Gainesville, FL USA
8	2- Department of Molecular Biosciences, Northwestern University, Evanston, IL USA
9	3- Department of Biology, Eastern Washington University, Cheney, WA USA
10	4- University of Florida Genetics Institute, Gainesville, FL USA
11	
12	Keywords: Caenorhabditis elegans, mutation accumulation, mutation rate, mutation spectrum,
13	private allele
14	

15 Abstract

16 Important clues about natural selection can be gleaned from discrepancies between the 17 properties of segregating genetic variants and of mutations accumulated experimentally under 18 minimal selection, provided the mutational process is the same in the lab as in nature. The ratio 19 of transitions to transversions (Ts/Tv) is consistently lower in C. elegans mutation accumulation 20 (MA) experiments than in nature, which has been argued to be in part due to increased 21 oxidative stress in the lab environment. Using whole-genome sequence data from a set of C. elegans MA lines carrying a mutation (mev-1) that increases the cellular titer of reactive oxygen 22 23 species (ROS), leading to increased endogenous oxidative stress, we find that the base-24 substitution spectrum is similar between *mev-1* lines, its wild-type progenitor (N2), and another 25 set of MA lines derived from a different wild strain (PB306). By contrast, the rate of short 26 insertions is greater in the *mev-1* lines, consistent with studies in other organisms in which 27 environmental stress led to an increase in the rate of insertion-deletion mutations. Further, the 28 mutational properties of mononucleotide repeats in all strains are qualitatively different from 29 those of non-mononucleotide sequence, both for indels and base-substitutions, and whereas 30 the non-mononucleotide spectra are fairly similar between MA lines and wild isolates, the 31 mononucleotide spectra are very different. The discrepancy in mutational spectra between lab 32 MA experiments and natural variation is likely due to a consistent (but unknown) effect of the lab 33 environment that manifests itself via different modes of mutability and/or repair at 34 mononucleotide loci.

35

36 Introduction

37 It is a fundamental principle of population genetics that the DNA sequence diversity in a population (θ) represents the product of mutation (μ) and "everything else", where "everything 38 39 else" subsumes the contributions of natural selection and random genetic drift in the composite 40 parameter N_e , the genetic effective population size: $\theta = 4N_e\mu$ (Watterson 1975; Nei and Li 1979). Empirically partitioning genetic variation into the contributions of mutation and everything else 41 42 requires that mutations be observed under conditions in which the effects of natural selection 43 are minimized as much as possible. That can be done in three basic ways: by genotyping 44 related individuals in a natural pedigree, as is now done commonly in humans (Gao et al. 2019; Halldorsson et al. 2019), by means of a "mutation accumulation" (MA) experiment, or by 45 cataloging very rare segregating variants that have (presumably) arisen only recently and thus 46 47 been minimally sieved by natural selection (Messer 2009; Carlson et al. 2018).

48 An MA experiment is in essence a large pedigree in which many replicate descendant 49 lines (MA lines) are derived simultaneously from a common ancestor and allowed to evolve 50 under conditions in which selection is minimal (Halligan and Keightley 2009). Typically, 51 selection in an MA experiment is minimized by minimizing N_{e} , ideally to a single individual or 52 chromosome. Pedigree genotyping has two important advantages over MA experiments: it can 53 be done in practically any organism, and it is fast. MA experiments are slow, labor intensive, 54 and are only practical in organisms with fast generation times that can be maintained easily in 55 the laboratory. However, MA experiments come with one unique advantage, which is that the 56 phenotypic effects of mutations can be assessed under controlled experimental conditions. MA 57 experiments have been a workhorse of evolutionary genetics for the past 60 years (Sprague et 58 al. 1960; Mukai 1964; Liu and Zhang 2019), and much of our understanding of the mutational 59 process has been derived from MA data (Drake et al. 1998; Halligan and Keightley 2009; Katju 60 and Bergthorsson 2019).

61 The utility of MA experiments is based on a key assumption: the mutational process under laboratory conditions faithfully reflects that in nature. It has long been known from studies 62 63 with microbes that all elements of the mutational process - rate, molecular spectrum, and phenotypic effects - depend on the environmental context and the genetic background; this 64 65 conclusion has recently been extended to multicellular eukaryotes (Sharp and Agrawal 2012; Sharp and Agrawal 2016; Kessler et al. 2020). On the one hand the context-dependence of the 66 67 mutational process is not a surprise; for example, the mutagenic effects of X-rays have been 68 known for a century. On the other hand, it suggests that MA experiments come with their own 69 biases (Baer 2019), just as natural selection biases the standing genetic variation.

70 The nematode *Caenorhabditis elegans* is an important model organism in many areas of 71 biology, and was the first metazoan organism to have its mutational process characterized at 72 the genomic level (Denver et al. 2000; Denver et al. 2004b). As genomic sequence data from 73 C. elegans MA lines and wild isolates have accumulated, it has become apparent that the base-74 substitution spectrum in lab-accumulated mutations differs from that of wild isolates in a 75 consistent way: there are more transversions in the lab than there are in nature. The ratio of 76 transitions to transversions (Ts/Tv) in MA lines is consistently around 0.6-0.8, whereas the 77 Ts/Tv ratio among wild isolates is around 1.1-1.2. Two early studies reported that $G:C \rightarrow T:A$ 78 transversions are overrepresented in MA data relative to the standing nucleotide diversity 79 (Denver et al. 2009; Denver et al. 2012); two more recent studies with more data reported that 80 A:T \rightarrow T:A transversions are also overrepresented in MA data relative to the standing nucleotide 81 diversity (Konrad et al. 2019; Saxena et al. 2019).

The MA spectrum may differ from the standing spectrum for several reasons. First, it may be that the mutational milieu in the lab is consistently different from that in nature. Second, it may be that purifying selection against transversions in nature is stronger than against transitions, and there is some reason to think that idea is plausible (Babbitt and Cotter 2011;

Guo et al. 2017). Third, it may be that the analyses of MA data and standing genetic variation
come with different biases.

One key difference between the lab MA environment and the natural environment is that 88 89 worms in the lab are fed ad libitum and kept at low population density and at a constant benign 90 temperature, which may result in a long-term elevation of metabolic rate relative to that 91 experienced in nature. A second key difference is that worms kept on plates in the lab 92 experience near-atmospheric concentrations of O₂, which is substantially greater than the 93 optimum, as assayed by worm preference (Gray et al. 2004). Both increased metabolic rate and 94 increased O₂ partial pressure could potentially increase the cellular concentrations of reactive 95 oxygen species (ROS). ROS are present in all cells as a natural byproduct of cellular metabolism, and can induce potentially mutagenic oxidative damage to DNA (Cooke et al. 2003; 96 97 Bridge et al. 2014) and alterations in chromatin structure (for review, Kreuz and Fischle 2016). 98 Accordingly, variation in cellular ROS levels has long been invoked as a potential cause of 99 variation in mutation rate (Richter et al. 1988; Shigenaga et al. 1989; Martin and Palumbi 1993; 100 Stoltzfus 2008). Cellular oxidative stress can vary due to differences in input (e.g., ROS levels 101 increase under conditions of physiological stress), in the ability of cells to enzymatically convert 102 ROS into benign products, and in repair processes (Beckman and Ames 1998; Turrens 2003; 103 Halliwell and Gutteridge 2007; Constantini 2014).

104 One well-documented manifestation of oxidative damage to DNA is the oxidation of 105 guanine to 8-oxo-7,8-dihydroguanine (8-oxodG), which if unrepaired, results in a $G \rightarrow T$ 106 transversion (Cheng et al. 1992; Cunningham 1997; Helbock et al. 1998; Cadet et al. 2003; 107 Evans and Cooke 2004). Accordingly, G:C \rightarrow T:A transversions are often interpreted as a signature of oxidative damage to DNA (Krašovec et al. 2017; Suzuki and Kamiya 2017; Poetsch 108 109 et al. 2018). While much of the early work linking ROS with 8-oxodG and mutation was 110 conducted in somatic cells (Dollé et al. 2000; Busuttil et al. 2003), this work has been extended to sperm (e.g., Paul et al. 2011; Kim and Velando 2020). Because previous studies have found 111

112 that the frequency of G:C \rightarrow T:A transversions is greater in MA lines than in the standing 113 variation, we were led to speculate that some property of the lab MA environment increases 114 oxidative stress relative to that experienced in nature. To test that hypothesis, we employed a 115 mutant strain of C. elegans, mev-1, that experiences elevated steady-state oxidative stress (Ishii 116 et al. 1998; Senoo-Matsuda et al. 2001; Ishii et al. 2013). We reasoned that, if $G:C \rightarrow T:A$ 117 transversions are in fact a signature of oxidative damage to DNA, then a strain with an 118 endogenously increased oxidative damage should show an even greater frequency of G:C→T:A 119 transversions than the background MA frequency. 120 Here we report results from an experiment in which a set of MA lines derived from a 121 strain carrying the mev-1(kn1) mutation backcrossed into our canonical wild-type N2 122 background were propagated for ~125 generations using our standard C. elegans MA protocol 123 (Joyner-Matos et al. 2011). We sequenced the genomes of 23 mev-1 lines and compared the 124 rate and spectrum of mutation to that of the N2 strain. As a further comparison, we report 125 results from an additional set of 67 MA lines derived from a different wild-type strain, PB306. 126 The laboratory MA spectrum is compared to the natural mutation spectrum inferred from "private 127 alleles" present as homozygous variants in one and only one wild isolate (n=773 wild isolates). 128 129 Results 130 Mutation Rate. 131 Mononucleotide repeats mutate differently than other sequence. We first report genome-wide

rates, which are ultimately of the most evolutionary relevance; we parse mutation rates into
 mononucleotide and non-mononucleotide rates in the next section. The experimental design is
 depicted in Figure 1.





Figure 1. Propagation of MA lines. The common ancestor of the MA lines (G0) was thawed from a cryopreserved sample and a single immature hermaphrodite picked onto an agar plate. Lines were propagated by single-worm transfer at four-day (one generation) intervals for t=Gmax transfers. Lines were initially genetically homogeneous (pink chromosome pairs);

t=Gmax transfers. Lines were initially genetically homogeneous (pink chromosome pairs); colored bars represent new mutations, which are fixed in a line with expectation u=(1-s)/(2-s).

where s is the selection coefficient (Keightley and Caballero 1997). See Methods and

141 where 5 is the selection coefficient (Keightey and Cabanero 1997). See Methods a

142 **Supplemental Table S1** for details of the MA experiments.

143 (i) Genome-wide mutation rate. Summary statistics on nuclear mutation rates are presented in

144 **Table 1** and results of statistical tests in **Supplemental Table S2**. Results for individual MA

145 lines are given in **Supplemental Table S3**. A complete list of mutations and their genomic

146 context is given in **Supplemental Table S4 (nuclear loci)** and **Supplemental Table S5**

147 (mtDNA loci). Raw sequence data are archived in the NCBI Short Read Archive, project

numbers PRJNA429972 (32 N2 MA lines) and PRJNA665851 (all other MA lines).

149 To test the hypothesis that the overall base-substitution mutation rate μ_{BS} differs

- between strains, we used a general linear model (GLM); details are presented in the Methods.
- 151 The hypothesis that the rate of G:C \rightarrow T:A transversions is greater in *mev-1* than in N2 is a prior
- one-tailed hypothesis test. The GLM revealed significant variation in μ_{BS} among the three

	<i>Mev-1</i> (<i>n</i> =23 MA lines)			N2 (<i>n</i> =68 MA lines)			PB306 (<i>n</i> =67 MA lines)		
	non-mono	mono	Total	non-mono	mono	Total	non-mono	mono	Total
$\mu_{GC \rightarrow AT}$	1.45 (0.12)	1.42 (0.36)	1.45 (0.10)	1.83 (0.06)	1.6 (0.24)	1.81 (0.06)	1.81 (0.05)	1.91 (0.22)	1.82 (0.05)
$\mu_{GC \rightarrow TA}$	0.81 (0.12)	1.29 (0.36)	0.85 (0.11)	0.80 (0.04)	1.13 (0.18)	0.82 (0.04)	0.94 (0.04)	1.68 (0.21)	1.00 (0.05)
µ _{GC→CG}	0.41 (0.06)	0.65 (0.26)	0.43 (0.06)	0.46 (0.04)	0.42 (0.09)	0.46 (0.03)	0.43 (0.03)	0.42 (0.09)	0.43 (0.03)
$\mu_{AT \to GC}$	0.39 (0.05)	0.33 (0.12)	0.38 (0.04)	0.47 (0.03)	0.40 (0.05)	0.46 (0.02)	0.45 (0.03)	0.34 (0.04)	0.43 (0.02)
$\mu_{AT \to CG}$	0.20 (0.03)	0.32 (0.10)	0.22 (0.03)	0.29 (0.02)	0.39 (0.05)	0.31 (0.02)	0.25 (0.02)	0.35 (0.04)	0.27 (0.02)
$\mu_{AT \to TA}$	0.35 (0.05)	3.04 (0.29)	0.83 (0.06)	0.44 (0.03)	3.34 (0.16)	0.95 (0.04)	0.40 (0.02)	2.70 (0.14)	0.81 (0.03)
μ _{BS}	1.60 (0.07)	3.62 (0.26)	1.89 (0.07)	1.91 (0.05)	3.93 (0.15)	2.20 (0.05)	1.89 (0.05)	3.52 (0.16)	2.13 (0.05)
μ _{INS}	0.15 (0.03)	1.32 (0.19)	0.32 (0.04)	0.07 (0.01)	1.07 (0.06)	0.21 (0.01)	0.10 (0.01)	1.19 (0.09)	0.26 (0.02)
μ _{DEL}	0.15 (0.02)	1.76 (0.23)	0.38 (0.03)	0.19 (0.01)	1.43 (0.09)	0.36 (0.02)	0.26 (0.02)	1.76 (0.12)	0.47 (0.03)
µ _{INDEL}	0.30 (0.05)	3.09 (0.22)	0.70 (0.04)	0.25 (0.01)	2.50 (0.11)	0.58 (0.02)	0.36 (0.02)	2.95 (0.17)	0.73 (0.03)
μτοται	1.90 (0.10)	6.70 (0.41)	2.59 (0.08)	2.17 (0.05)	6.43 (0.20)	2.78 (0.06)	2.25 (0.06)	6.47 (0.22)	2.85 (0.07)
U _{GENOME}	0.16	0.10	0.25	0.18	0.09	0.27	0.19	0.09	0.27

Table 1. Mutation rates (μ) x 10⁹/site/generation (SEM). U_{GENOME} is the haploid genome-wide mutation rate per-generation.

strains, as well as a significant interaction between strain and mutation type. Contrary to our expectation, μ_{BS} summed over all six mutation types is significantly less in the *mev-1* lines than in N2, and marginally lower than in PB306. In contrast, μ_{BS} does not differ between N2 and PB306. The G:C \rightarrow T:A transversion rate differs among strains (**Figure 2A**), but not in the predicted way: $\mu_{GC\rightarrow TA}$ is indistinguishable between *mev-1* and N2, and lower than in PB306. The rate of A:T \rightarrow T:A transversions is marginally different between the three strains (**Figure 2A**), with N2 having a slightly greater rate than *mev-1* and PB306.



163

Figure 2. Mutation rates. Left (A-C), type-specific base-substitution rates, by strain. Right (D-F), size-specific indel rates, by strain. Top row (A,D), genome-wide mutation rates; middle row (B,E), rates at non-mononucleotide repeat sequence; bottom row (C,F), mononucleotide repeat sequence. All rates are scaled as 10^{-9} per base per generation; note the difference in y-axis scale between panels (A-C, left) and panels (D-F, right). Error bars represent 1 SEM.

169 The insertion and deletion rates both vary among strains, in different ways (**Figure 2D**;

170 **Table 1**). The PB306 deletion rate is significantly greater than that of *mev-1* and N2. The *mev-1*

insertion rate is significantly greater than that of N2 and greater than that of PB306, although not
 significantly. The PB306 insertion rate is marginally greater than that of N2.

173 The mechanisms of mutagenesis and repair are not necessarily uncorrelated, and the 174 large number of mutations (>9,000) provides an opportunity to investigate the covariance structure of the different types of mutation. We first investigated the covariance structure of the 175 176 six types of base-substitution. We initially fit a model with a single variance component, i.e., a 177 uniform diagonal element in the covariance matrix and all off-diagonal elements constrained to 178 zero. That model was compared to a model in which a separate variance (diagonal element) 179 was estimated for each of the six traits ("banded main diagonal") and the off-diagonal elements 180 constrained to zero. That model was in turn compared to a model in which all elements are unconstrained. The unconstrained model had the smallest AICc (Δ AICc=16.2 less than the 181 182 banded main diagonal), and provided a significantly better fit to the data (LRT, -2Δ InL=47.1, 183 df=15, P<0.0001), although the individual correlations were small (all |r|<0.3; Supplemental 184 Table S6). A model with the covariance matrix estimated separately for each strain did not fit 185 as well (Δ AICc=12.4 greater than the model with a single covariance matrix).

186 We next investigated the covariance structure of indels and base-substitutions together, 187 combining all six base-substitutions into one category with insertions and deletions included 188 separately. The best model is the most parameter-rich, with the unstructured covariance matrix 189 estimated separately for each strain (Δ AlCc=12.3 less than the model with a single unstructured 190 covariance matrix; LRT, -2Δ InL=37.6, df=12, P<0.0002). In all three strains, correlations 191 between base-substitution and deletion rates are moderately positive (r=0.2-0.3), and base-192 substitution and insertion rates are uncorrelated (r≈0). The correlation between base-193 substitution rate and insertion rate is negative in *mev-1* and N2, whereas it is moderately 194 positive in PB306 ($r \approx 0.3$; Supplemental Table S7). We emphasize that, for both of these 195 covariance analyses, we cannot reject the hypothesis that any specific off-diagonal element is different from zero. Rather, the results simply mean that models with some non-zero off 196

diagonal elements fit the data better than a model in which all off-diagonal elements are
constrained to zero, and we interpret the point estimates of the individual pairwise correlations
heuristically.

200 Inspection of the residuals of the GLM reveals two MA lines that are obvious outliers. 201 Line 572 (N2) has anomalously high base-substitution and deletion rates, although its insertion 202 rate is slightly below the N2 average. Line 481 (PB306) has anomalously high insertion and 203 deletion rates, although its base-substitution rate is only slightly greater than the PB306 average (Supplemental Table S3). We searched the list of mutations in those two lines for candidates 204 205 that could potentially increase the mutation rate (see Extended Methods, Supplemental 206 Appendix A1.4). It turns out there are obvious candidates in both lines (Supplemental Table 207 **S4**). Line 572 has a T \rightarrow G transversion upstream of the *atl-1* gene; *atl-1* is involved in 208 homologous recombination and the DNA damage checkpoint. Line 471 has a 1-bp deletion 209 downstream of the xpc-1 gene; xpc-1 is involved in nucleotide excision repair.

210 The mitochondrion is the site of the electron transport chain subunit II, and therefore the 211 source location of the excess free radicals produced in *mev-1* worms (Senoo-Matsuda et al. 212 2001). However, the mtDNA mutation rate μ_{Mt} is not increased in *mev-1*, and is very close to that of N2 (*mev-1* μ_{Mt} = 5.62+2.62x10⁻⁸/gen; N2 μ_{Mt} = 6.05+1.36x10⁻⁸/gen; randomization test 213 p>0.86). The point estimate of the mtDNA mutation rate is ~50% greater in PB306 (μ_{Mt} 214 = $8.84+2.15\times10^{-8}$ /gen), but the difference between PB306 and N2 is not significantly different 215 from zero (randomization test p>0.47). A list of mtDNA mutations and their heteroplasmic 216 217 frequencies is given in **Supplemental Table S5**. Details of the calculation of μ_{Mt} and of the 218 randomization test are given in the Methods.

(ii) Mononucleotide repeats. Mononucleotide repeats are well-known to incur indel mutations at
 a much greater rate than non-repeat sequence (Denver et al. 2004a), and we previously found
 that mononucleotide repeats in the N2 genome of eight or more consecutive bases also
 experience an elevated rate of base-substitution mutations (Saxena et al. 2019). Here we find

223 the same result when mononucleotides are defined by the less stringent criterion of ≥5 224 consecutive bases. Mononucleotide repeats of ≥5 bases comprise ~8.5% of the N2 genome, of 225 which ~80% are A:T repeats (see Extended Methods, Supplemental Appendix A1.8). 226 Pooled over strains and mutation types, the mononucleotide base-substitution rate is nearly 227 twice that of non-mononucleotide sequence (**Table 1**). However, the increased base-228 substitution mutation rate in mononucleotide repeats is not uniform; the rate of transversions 229 from either a G:C or an A:T base pair to an A:T base pair is greater in mononucleotide repeats, 230 whereas rates of the other four types of base-substitution do not differ between sequence types (Figure 2B,C). In particular, the rate of A:T \rightarrow T:A transversions is approximately seven-fold 231 greater in mononucleotide repeats than in non-repeat sequence. 232 As expected, the rate of +/- 1 bp indel mutations is much greater in mononucleotide 233 234 repeats than in non-repeat sequence; averaged over strains, 1 bp deletions occur about 26 235 times more frequently and 1 bp insertions about 39 times more frequently in mononucleotides 236 than in non-mononucleotide sequence. In contrast, the rate of both deletions and insertions 237 longer than 1 bp is only about twice as great in mononucleotides as in non-mononucleotide 238 sequence (Figure 2D-F; Table 1). Summed over sequence types and mutation types, the rate of 1 bp indels is about 25% greater in mev-1 than in N2, whereas the rates in N2 and PB306 are 239 240 similar.

241

242 Local sequence context.

A previous analysis of the factors potentially affecting base-substitution mutability in N2 revealed a predominant role for the local (three-nucleotide) sequence context (Saxena et al. 2019). In that study, motifs with C or G in the mutant (3') site were more mutable on average than motifs with A or T in the mutant site, with one exception: the triplet 5'-ttA-3' was the most mutable of any of the 64 motifs. Subsequent MA studies with the N2 strain of *C. elegans* have found a similarly high mutability of the 5'-ttA-3' motif (Konrad et al. 2019; Volkova et al. 2020).

The 64 nucleotide triplets are depicted in **Figure 3**, oriented $5' \rightarrow 3'$ with the mutant site in the 3' 249 position. There are not sufficiently many mutations to permit a robust statistical comparison 250 251 between strains of the full set of 64 triplets, but we have a prior hypothesis with respect to the 252 relative mutability of the 5'-ttA-3' triplet. 5'-ttA-3' is also the most mutable motif in the other two 253 strains, although the size of the anomaly is not as extreme in the other strains as in N2. The 5'ttA-3' mutation rate differs among strains (N2, μ_{ttA} =8.6x10⁻⁹/gen; mev-1, μ_{ttA} =6.6 x10⁻⁹/gen; 254 PB306, μ_{ttA} =6.4x10⁻⁹/gen); the difference between PB306 and N2 is statistically significant. The 255 256 correlation of the 64 motif-specific mutation rates between strains is large and positive (>0.7). 257 but significantly less than 1 in all three cases (Supplemental Table S8). 258 Some fraction of 5'-ttA-3' triplets occur at the 3' end of poly-T repeats, suggesting the 259 possibility that the atypically high mutability of that motif results from the association of the motif 260 with mononucleotide repeats. Averaged over the three strains, μ_{ttA} when the motif is adjacent to 261 a poly-T repeat is ~27-fold greater than when the same motif is not adjacent to a poly-T repeat $(43.0 \pm 2.3 \times 10^{-9})$ /generation vs. $1.6 \pm 0.2 \times 10^{-9}$ /generation). When not in the context of a 262 263 mononucleotide repeat, the 5'-ttA-3' motif mutates at essentially the same rate as any A:T base-264 pair (Figure 3). Details of the local sequence context analysis are presented in the **Extended Methods**, 265

266 Supplemental Appendix A1.9.

267



268

Figure 3. 64 3-bp motif base-substitution mutation rates. Motifs are arranged 5'-xyZ-3', with the mutant base Z in the 3' position. Error bars show one SEM.

- 272 Mutation Spectrum.
- 273 The mutation spectrum, defined as the frequency distribution of mutations of given types, is not
- the same as the distribution of type-specific mutation rates, although the two are obviously
- related. In our MA data, pooling over strains and the six base-substitution types, the correlation
- between the type-specific base-substitution mutation rate (μ_i) and the proportion of mutations of

that type (p_i), $r_{\mu,p}$ = 0.85. The utility of knowing the mutation spectrum in MA lines is that it provides a benchmark for comparison to wild isolates, for which the mutation rate cannot be known.

Summed over all MA lines within strains, the six-category base-substitution mutation spectrum of *mev-1* (*n*=508 mutations) does not differ from those of either N2 (*n*=3,434 mutations; 2x6 Monte Carlo Fisher's Exact Test, 10⁶ iterations, P>0.50) or PB306 (*n*=3,111 mutations; MC FET, P>0.45), whereas the spectrum of N2 differs significantly from that of PB306 (MC FET, P<0.001; **Figure 4A**). The differences between the base-substitution spectra of N2 and PB306 are not great, but the large number of mutations enables us to detect significant differences on the order of a few percent.



Figure 4. Base-substitution spectra. Left panel (A-C), MA lines; Right panel (D-F), wild isolate private alleles (blue) and MA means (red). Top panels (A,D), whole-genome; middle

panels (B,E), non-mononucleotide sequence; bottom panels (C,F), mononucleotide sequence.
 Error bars show one SEM.

293	To compare the indel spectra between strains, we first investigated the overall mutational bias,
294	quantified as the proportion of deletions among indel mutations. The indel bias differed
295	significantly among strains (3x2 Fisher's exact test, P<0.04). Breaking down the bias into
296	pairwise comparisons between strains, mev-1 (55% deletion) had a significantly lower
297	proportion of deletions than N2 (63% deletion, Fisher's Exact Test P< 0.04) and PB306 (65%
298	deletion, FET P<0.02); the bias did not differ significantly between N2 and PB306.
299	To characterize the indel spectra with finer resolution, we next assigned insertions and
300	deletions separately to four bins of size 1, 2-5, 6-10 and 11-20. The bin sizes are obviously
301	arbitrary, but the distribution provides cell counts of five or more observations for all but the
302	largest insertion bin in the mev-1 strain. The genome-wide spectrum (Figure 5A) did not differ
303	between <i>mev-1</i> and N2 (FET, p>0.08) nor between N2 and PB306 (FET, p>0.29).



304

Figure 5. Indel spectra. Left panel (A-C), MA lines; Right panel (D-F), wild isolate private
 alleles (blue) and MA means (red). Top panels (A,D), whole-genome; middle panels (B,E), non mononucleotide sequence; bottom panels (C,F), mononucleotide sequence. Error bars show one
 SEM.

- 310 Breaking the genomic context into mononucleotide repeats and non-mononucleotides, the non-
- mononucleotide spectra (Figure 5B) were marginally different between mev-1 and N2 (FET,
- p<0.03), but not between N2 and PB306 (FET, p>0.11). The mononucleotide spectra (Figure
- **5C**) were very similar between *mev-1* and N2 (FET, p>0.82), but significantly different between
- N2 and PB306 (FET, p<0.01). Averaged over N2 and PB306, the deletion bias is stronger in
- non-mononucleotide sequence (72.5% deletion) vs. mononucleotides (58% deletion).
- 316
- 317 Comparison of the MA spectrum to the natural private allele spectrum.

318 (i) Base-substitutions. Variant alleles present in a single wild isolate ("private alleles")

319 presumably arose as new mutations in the recent past, and have been minimally scrutinized by 320 selection. As expected from previous studies, the genome-wide base-substitution spectrum of 321 unique variants is qualitatively different from those of the MA lines (Figure 4B). Because of the 322 large sample sizes, even small differences in proportions between groups are highly statistically 323 significant (FET, P<<0.0001 in all cases), so we interpret the differences heuristically. We first 324 consider the genome-wide spectrum, then parse the genome into mononucleotide repeats and 325 non-mononucleotide repeats. The wild isolate spectra are compared to the average of the N2 326 and PB306 MA spectra (which are themselves significantly different). The full CeNDR wildisolate dataset includes 773 unique genotypes, which collectively harbor >450,000 private 327 single-nucleotide variants. Of those 773 wild isolates, we arbitrarily restricted our analysis to 328 329 447 isolates which carry 100 or fewer private alleles (n=11,667 private alleles), which represent 330 about 400 generations of mutation accumulation. A list of the wild isolates included and the set of private alleles for each isolate are given in Supplemental Table S9. 331

A:T \rightarrow T:A transversions occur more frequently in the MA lines than in the wild isolates, and A:T \rightarrow G:C transitions occur less frequently in the MA lines. G:C \rightarrow T:A transversions, the original focus of this study, occur only slightly more frequently in the MA lines than among the rare variants. The wild isolate private allele spectrum is closer to that of PB306 (and *mev-1*) than it is to that of N2, but the MA spectra are clearly more similar to each other than either is to the wild isolate spectrum.

Comparison of the base-substitution spectra of mononucleotide repeat sequences to the spectra of non-mononucleotide sequences is revealing (**Figure 4B**). For non-mononucleotide sequence (~90% of the genome), the private allele spectrum of wild isolates is similar to that of the MA lines, although the proportion of private alleles mutating from a C:G base-pair is less than the MA frequency in all three cases. In contrast, the mononucleotide spectra are very different. The proportion of A:T \rightarrow T:A transversions in the wild isolates is less than half that

expected from the MA proportion (28% vs. 65%), and there are many fewer A:T \rightarrow G:C

transitions in the MA lines than in the private alleles (8% vs. 25%)

346 If the strength of purifying selection varies among the different types of mutations, we 347 expect the spectrum of common alleles will differ from that of rare variants. We compared the 348 genome-wide allele frequency spectrum between private alleles and common(er) alleles. 349 defined as variants present in more than one isolate but with an upper bound of 10% on the 350 minor allele frequency. We chose 10% as the upper-bound on the minor allele frequency to 351 minimize the chance of misidentifying an ancestral allele as the recent mutant. The spectra of 352 rare and common variants are similar (Supplemental Figures S1, S2), reinforcing the notion that the difference in spectra between standing variants and mutations accumulated in 353 354 laboratory MA experiments results from a consistent difference between the mutational milieu in 355 the lab and in nature, rather than from the action of differential purifying selection.

356 As noted, the MA spectrum differs slightly but significantly between N2 and PB306. N2 357 is long-adapted to the lab environment, and differs from wild-type C. elegans in many respects 358 (Sterken et al. 2015), of which the mutation spectrum is but one. Private alleles in the wild 359 isolates provide a much broader sample of the mutational process in the species. Parametric 360 bootstrap simulations (see Methods for the details) show that there is more variation in the 361 spectrum of private alleles among wild isolates than expected from a uniform base-substitution 362 mutational process (Supplemental Figure S3), both for non-mononucleotide sequence and mononucleotide runs, as well as genome-wide. This finding demonstrates that, unsurprisingly, 363 364 the C. elegans mutational process varies among strains in nature as well as in the lab, although the causes of the variation in nature are uncertain. 365

<u>(ii) Indels</u>. The frequencies of false positives and false negatives are qualitatively greater for
 indels of >20 bp than for smaller indels (see next section), so we restrict comparison of indel
 spectra of MA lines and wild isolates to indels of <21 bp. To compare the indel spectra of MA

lines and natural private alleles we applied additional filters on the MA lines and used the full set

370 of 773 wild isolates (Extended Methods, Supplemental Appendix A1.11).

371 As noted previously, the MA lines have a strong genome-wide deletion bias (~62% 372 deletion); the deletion bias in the wild isolates is much weaker (54% deletion). Inspection of the 373 genome-wide spectrum (Figure 5D) reveals that +/- 1 bp indels are much a much greater 374 fraction of indels in the MA lines than in the wild isolates. Breaking the genome into 375 mononucleotide and non-mononucleotide components, it is apparent that the non-376 mononucleotide spectra are roughly congruent between MA lines and wild isolates (Figure 5E). 377 In contrast, the mononucleotide spectra (Figure 5F) are different between MA and wild isolates, with the MA lines having a greater fraction of \pm 1 bp indels. Evidently, the genome-wide 378 379 discrepancy between the MA indel spectrum and the private allele spectrum of wild isolates is 380 largely due to the different mutational properties of mononucleotide repeats in lab populations 381 and in nature. 382 False positives and false negatives 383 For an estimate of mutation rate to be believable, it is necessary to have credible estimates of

the frequency of false positives (FP, apparent variants called as new mutations that are not new mutations) and false negatives (FN, mutated sites in the genome that are not called as new mutations). We designate alleles that are homozygous in the *C. elegans* reference genome (N2) as "0"; variant alleles are designated as "1". Some sites in the common ancestors of our MA lines differ from the reference allele, i.e., the MA ancestor genotype is 1/1 (1,439 sites in the N2 ancestor, 1,642 in the *mev-1* ancestor, 155,614 in the PB306 ancestor).

False positives were assessed in two ways. First, we counted sites that were scored 1/1 in our MA ancestor and 0/0 in the reference genome. These are presumed to be new mutations fixed subsequent to the divergence of the MA ancestor and the reference strain from their common ancestor (alternatively, they could be errors in the reference genome). Then, we scored those sites in each MA line in the set. If a site was scored 1/1 in the MA ancestor but

scored 0/0 in all MA lines, we inferred a false positive in the MA ancestor. By that criterion, the
false discovery rate (FDR=FP/[FP+TP]) in the N2 MA lines for base-substitution mutations is
0.19% and in PB306 it is 0.0033%. FDRs for indels in the two strains are 0 and 0.28%,
respectively.

399 Second, we employed an independent set of low-coverage sequence data from a set of 400 192 recombinant inbred advanced intercross lines (RIAILs) generated from a cross between two 401 N2 strain MA lines, 530 and 563. The details of the construction and sequencing of the RIAILs and data analysis are presented in **Supplemental Appendix A2**: RIAIL genotypes are given in 402 403 Supplemental Table S10. Each parental line carries its own set of putative mutations, which are expected to segregate in the RIAILs with an average frequency of 50%. A variant called as 404 405 a mutant in a parental line (1/1) that does not segregate in the RIAILs is designated as a false 406 positive in the parent. The false discovery rate for detection of SNPs and indels is 2.1% and 407 2.9%, respectively. This result is consistent with a previous estimate of the FDR from N2 strain 408 MA lines based on Sanger sequencing of PCR products, which gave an upper 95% confidence 409 limit on the FDR of 2.5% (Saxena et al. 2019).

True mutations may not be identified as such for two reasons. First, the mutant site may not be covered by the sequencing. We refer to this situation as "failure to recall" a mutation. Or, a true mutation (i.e., 0/0 in the ancestor and 1/1 in the MA line carrying the mutation) may be incorrectly called as "not 1/1", i.e., either as 0/0 or as a heterozygote, 0/1. These are false negatives. False negatives will cause the mutation rate to be underestimated. Failure to recall mutations will only cause the mutation rate to be misestimated (under or over) if the mutation rate at sites not covered in the sequencing differs from that of the sites covered.

We employed a simulation approach to assess recall rate, the details of which are given in the **Extended Methods, Supplemental Appendix A1.10**. "Dummy" mutations were introduced into the reference genome at random, and the MA data analyzed using our standard variant-calling pipeline but with the simulated genome as the reference. Sites with dummy

mutations are scored as 1/1 in the MA ancestor and all MA lines. If a dummy mutation at a site was not called 1/1 in an MA line, for any reason, it was classified as a failure to recall. The genome-wide base-substitution failure-to-recall rate is 6.81% in N2, 6.92% in *mev-1*, and 9.18% in PB306. The small indel (\leq 50bp) failure-to-recall rates are 9.34% in N2, 9.78% in *mev-1*, and 12.35% in PB306.

Dummy mutations that are called as "not 1/1" (i.e., 0/0 or 0/1) are false negatives *sensu* stricto. The genome-wide base-substitution false negative rate (FNR) is 0.12% in N2, 0.17% in mev-1, and 0.91% in PB306. The small indel (\leq 50bp) FNRs are 5.15% in N2, 5.59% in mev-1, and 5.84% in PB306.

430

431 Discussion

432 Four key findings emerge. First, the base-substitution mutational process of *mev-1* is similar to 433 that of N2, from which it was derived. Therefore, any discrepancy between the basesubstitution spectrum of N2 and those of wild isolates is unlikely to be due to differences 434 435 resulting from endogenous oxidative stress manifested in the lab environment. Moreover, the 436 overall base-substitution rate of mev-1 is significantly less (~14%) than that of N2, contrary to 437 the prediction that elevated levels of ROS increase the base-substitution mutation rate. In 438 contrast, the indel rate, and especially the insertion rate, is greater in the *mev-1* lines than in N2 439 (~50% greater). We have previously shown that the dinucleotide repeat indel rate increases in N2 MA lines propagated under conditions of heat stress (Matsuba et al. 2013), and a similar 440 increase in the indel rate - but not the base-substitution rate - has been shown in Arabidopsis 441 thaliana MA lines cultured under conditions of heat stress (Belfield et al. 2020). Drosophila 442 melanogaster MA lines initiated from genomes carrying deleterious large-effect mutations 443 444 experience higher rates of short deletions than wild-type MA lines, thought to be due to 445 preferential deployment of different mechanisms of double-strand break repair (Sharp and Agrawal 2016). The increased insertion rate in the mev-1 lines suggests that elevated levels of 446

endogenous oxidative damage may in fact influence the mutational process, but not in the waythat we predicted.

449 A possible explanation for the lower base-substitution rate in *mev-1* relative to N2 is that 450 the N2 lines underwent approximately twice as many generations of MA (Gmax=250 451 generations) as did *mev-1* (Gmax=125 generations). We previously found that "second order" 452 N2 MA lines initiated from a subset of the N2 lines included in this study evolved a significantly 453 greater (~10%) mutation rate over a subsequent ~150 generations of MA. The results reported 454 here are consistent with the N2 (and PB306) lines evolving an increased base-substitution 455 mutation rate over time, although the evidence is circumstantial. The second key result emerges from the comparison of the mutational properties of the 456 457 lab-adapted N2 strain and the wild isolate PB306. The overall base-substitution mutation rates 458 are very similar between the two strains (<5% different), although some of the type-specific 459 mutation rates are marginally different (e.g., the genome-wide G:C \rightarrow A:T transition rate is ~20% 460 greater in PB306, and the A:T \rightarrow T:A transversion rate is ~20% greater in N2). The type-specific 461 differences between strains are not large – by way of contrast, two sets of *D. melanogaster* MA 462 lines derived from different starting genotypes had a five-fold difference in $G:C \rightarrow A:T$ transition 463 rates (Schrider et al. 2013). PB306 has a similarly greater indel rate than N2 (~25%), although 464 the direction and magnitude of the bias are nearly identical (65% and 63% deletion,

465 respectively).

The differences in mutational properties between N2 and PB306 surely have a genetic basis. The two strains' genomes are about 1.5% different (~150K pairwise differences). With only two strains, we cannot say whether the difference is related to the domestication of N2, although it is interesting that the mutational properties of *mev-1* trend in the same direction as those of PB306. If the elevated endogenous oxidative stress experienced by *mev-1* is reflective of environmental stress in general, it suggests that the lab environment may be slightly stressful for PB306 compared to the lab-adapted N2. Moreover, the private allele spectrum of the wild

473 isolates is closer to that of PB306 than of N2, which, by the preceding logic, suggests that the lab environment may be especially comfortable for N2 relative to the natural environment. The 474 475 finding that there is too much variation in the standing mutation spectrum among wild isolates to 476 be explained by a uniform mutational process suggests that the genetic variation in the 477 mutational process observed between N2 and PB306 is not an artifact of the lab environment. 478 Probably the most important point to take away from the comparison, however, is that both the 479 rate and spectra of N2 and PB306 are not very different, which implies that conclusions about 480 mutation drawn from N2 MA lines may be generalizable to *C. elegans* at large.

481 The third key finding is the discrepancy in the indel bias between the MA lines and the wild isolate private alleles: MA lines show a strong deletion bias (~62% genome-wide), whereas 482 the deletion bias in the private alleles is much weaker (~54% deletion). As noted above, the 483 484 indel rate has been shown to increase under stress in a variety of experimental systems, 485 including C. elegans. The reduced deletion bias in mev-1 relative to N2 reinforces the inference 486 that a nematode's life in nature is more stressful than life in the lab, with the attendant 487 differences in the mutational process. To the extent that mutational variance in phenotypic traits 488 is underlain by indel mutations, evolutionary inferences that rely on comparisons between 489 mutational and standing genetic variance are rendered less robust.

490 The fourth key finding is the qualitative difference in the base-substitution mutation 491 spectra between mononucleotide repeat sequence and non-mononucleotide sequence in the 492 MA lines, and how the MA spectra relate to the standing private allele spectra. This study was 493 predicated on the discrepancy between the (N2) MA base-substitution spectrum and that seen 494 in wild isolates. In fact, when the non-mononucleotide spectra are compared, the discrepancy 495 substantially disappears. Although significant differences remain, they do not appear wildly 496 different (Figure 4). Conversely, there is a huge discrepancy between the mononucleotide 497 spectra of MA lines and private alleles, and in particular with respect to how A:T mutates. First, a larger fraction of mutations occur at A:T sites in the MA lines than in the private alleles (81% 498

499 vs. 66%). Second, there are many more A:T \rightarrow T:A transversions in the MA lines than in the 500 private alleles (65% vs. 28%) and there are many fewer A:T \rightarrow G:C transitions in the MA lines 501 than in the private alleles (8% vs. 25%). The technology employed in sequencing most (not all) 502 of the wild isolates and most (not all) of the MA lines was the same, and the data were analyzed 503 in the same way (Extended Methods, Supplemental Appendix A1.4; Supplemental Figure 504 S4), so it is difficult to imagine that the discrepancy is due to wildly different rates of false 505 positives or false negatives in one or the other set of data. Given that the genetic basis of 506 eukarvotic DNA repair is complex, and hundreds of genes are known to influence variation in 507 DNA repair (Eisen and Hanawalt 1999), a reasonable guess is that some element(s) of DNA 508 repair differs systematically between the lab environment and the natural environment, but what 509 it may be we cannot say.

510

511 Methods and Materials

512 Strains.

513 (i) MA lines. The *mev-1* gene encodes a subunit of succinate dehydrogenase cytochrome b, a 514 component of complex II of the mitochondrial electron transport chain. The mev-1(kn1) allele is 515 a $G \rightarrow A$ transition that replaces a glycine with a glutamate, resulting in decreased enzyme 516 activity and increased electron leak. The resultant phenotype has been explored in several 517 species (Ishii et al. 1990; Ishii et al. 2011; Ishii et al. 2013; Ishii et al. 2016) and includes elevated 8-oxodG and increased nuclear mutation rate (Hartman et al. 2004; Ishii et al. 2005). 518 519 The mev-1(kn1) allele was backcrossed into the canonical Baer lab N2 strain genomic 520 background; see Extended Methods, Supplemental Appendix A1.1 for details. PB306 is a wild isolate generously provided by Scott Baird; the N2 lines are derived from 521 522 a replicate of the ancestor of the Vassilieva and Lynch (1999) MA lines.

- 523 (ii) Wild isolates. Genome sequence data for 773 unique wild isolates were obtained from the *C*.
- 524 *elegans* Natural Diversity Resource (CeNDR; Cook et al. 2017). Details of the sequencing and
- 525 variant calling are available at https://www.elegansvariation.org/.
- 526

527 *Mutation accumulation experiments.*

- 528 Details of the N2 and PB306 mutation accumulation experiment are given in Baer et al. (2005),
- those of the *mev-1* MA experiment are given in Joyner-Matos et al. (2011), and are summarized
- 530 in the **Extended Methods**, **Supplemental Appendix A1.2**. Data on MA line transfers is
- presented in **Supplemental Table S1**. *mev-1* MA lines were propagated for up to 125
- 532 generations of MA (Gmax=125); N2 and PB306 lines were propagated for up to 250
- generations of MA (Gmax=250). The basic protocol in both experiments follows that of

534 Vassilieva and Lynch (1999) and is depicted in **Figure 1**.

- 535
- 536 Genome sequencing and bioinformatics

537 32 N2 lines and 20 PB306 lines were previously sequenced with Illumina short-read sequencing 538 at an average coverage of ~ 27X read-depth with 100 bp paired-end reads. The N2 data have 539 been previously reported (Saxena et al. 2019). For this project we sequenced an additional 40 540 N2 lines, 54 PB306 lines, and 24 mev-1 lines and their ancestors using Illumina short-read 541 sequencing at an average coverage of 46X depth with 150 bp paired-end reads. Protocols for 542 DNA extraction and construction of sequencing libraries are given in the **Extended Methods**, 543 Supplemental Appendix A1.3; details of bioinformatics processing of raw sequence data and 544 variant calling are given in the Extended Methods, Supplemental Appendix A1.4. Following preliminary analysis, variants were called using HaplotypeCaller (in 545 546 BP_RESOLUTION mode) in GATK4 (v4.1.4.0) (McKenna et al. 2010). Variants were identified 547 as putative mutations if (1) the variant was identified as homozygous, and (2) it was present in

one and only one MA line. Criterion (1) means that any mutations that occurred in the last few

549 generations of MA that were still segregating and/or occurred during population expansion for DNA extraction were ignored. Because the MA progenitor was at mutation-drift equilibrium 550 551 (Lynch and Hill 1986), the segregating variation is expected to be the same in the MA progenitor 552 and the MA line, so ignoring heterozygotes results in an unbiased estimate of mutation rate. 553 Criterion (2) reduces the probability of mistakenly identifying a variant segregating at low 554 frequency in the expanded population of the MA progenitor as a new mutation. One mev-1 line 555 was found not to carry the *mev-1(kn1)* allele and was omitted as a presumed contaminant. Four 556 pairs of N2 strain MA lines and seven pairs of PB306 strain lines shared multiple variants and 557 were inferred to have experienced contamination at some point during the MA phase; one line from each pair was arbitrarily omitted from subsequent analyses. The final data set includes 68 558 559 N2 lines, 67 PB306 lines, and 23 mev-1 lines (Supplemental Table S3).

560

561 Data Analysis.

562 (i) Nuclear mutation rate. Because different lines experienced different numbers of generations 563 during the MA phase, and because the number of callable sites differs among lines, we cannot 564 use the number of mutations per line to directly compare mutation rates among strains. Instead, 565 we calculated a mutation rate μ_i for each line i as $\mu_i = m_i / n_i t_i$, where m is the number of mutations, 566 *n* is the number of callable sites, and *t* is the number of generations of MA (Denver et al. 2009). 567 To test for group-specific effects on mutation rate, and to partition the (co)variance in mutation rate into within and among-group components, we consider the mutation rate itself as a 568 569 continuously-distributed dependent variable in a general linear model (GLM). Details of the 570 GLM analyses are given in the Extended Methods, Supplemental Appendix A1.5. 571 (ii) mtDNA mutation rate. Estimation of the mtDNA mutation rate is more complicated than for 572 nuclear loci because a non-trivial fraction of mutations will not have reached fixation and remain 573 heteroplasmic. The probability of a mutation ultimately reaching fixation is its current frequency 574 in the population (Wright 1931), so the mtDNA mutation rate of MA line *i* can be estimated as

575 $\mu_i = \frac{\sum p_j}{n_i t_i}$, where p_j is the frequency of the *j*th mutation in line *i*, n_j is the number of callable

- 576 sites in line *i*, and *t_i* is the number of generations of MA of line *i* (Konrad et al. 2017). However,
- 577 since many lines have no mtDNA mutations, GLM is not an appropriate analysis. Instead, we
- used randomization tests of hypothesized differences in mtDNA mutation rate; the details are
- 579 presented in the Extended Methods, Supplemental Appendix A1.6
- 580 (iii) Mutation spectrum. Mutation spectra were compared between groups by Fisher's Exact
- 581 Test. For comparisons with sample sizes too large to calculate directly we used Monte Carlo
- estimates of the FET as implemented in the FREQ procedure of SAS v.9.4.
- 583 We also tested the hypothesis that the variance in the private allele frequency spectrum
- among wild isolates can be explained by sampling variance around a single uniform base-
- substitution spectrum with expectation equal to the observed frequencies. We employed
- 586 parametric bootstrap simulations, the details of which are presented in the **Extended Methods**,
- 587 Supplemental Appendix A1.7.
- 588

589 Acknowledgments

- 590 We thank J. Dembek for laboratory assistance. Support was provided by NIH awards
- 591 GM107227 to CFB and ECA, and GM127433 to CFB and V. Katju.

592 References Cited

- Babbitt GA, Cotter CR. 2011. Functional conservation of nucleosome formation selectively biases
 presumably neutral molecular variation in yeast genomes. *Genome Biology and Evolution* 3: 15 22.
- Baer CF. 2019. Evolution: Environmental dependence of the mutational process. *Current Biology* 29(11):
 R415-R417.
- Baer CF, Shaw F, Steding C, Baumgartner M, Hawkins A, Houppert A, Mason N, Reed M, Simonelic K,
 Woodard W et al. 2005. Comparative evolutionary genetics of spontaneous mutations affecting
 fitness in rhabditid nematodes. *Proceedings of the National Academy of Sciences of the United* States of America 102(16): 5785-5790.
- 602 Beckman KB, Ames BN. 1998. The free radical theory of aging matures. *Physiol Rev* 78(2): 547-581.
- 603Belfield EJ, Brown C, Ding ZJ, Chapman L, Luo M, Hinde E, van Es SW, Johnson S, Ning Y, Zheng SJ et al.6042020. Thermal stress accelerates Arabidopsis thaliana mutation rate. Genome Research.
- 605Bridge G, Rashid S, Martin SA. 2014. DNA mismatch repair and oxidative DNA damage: Implications for606cancer biology and treatment. Cancers 6: 1597-1614.
- Busuttil RA, Rubio M, Dollé MET, Campisi J, Vijg J. 2003. Oxygen accelerates the accumulation of
 mutations during the senescence and immortalization of murine cells in culture. Aging Cell 2:
 287-294.
- Cadet J, Douki T, Gasparutto D, Ravanat J-L. 2003. Oxidative damage to DNA: formation, measurement
 and biochemical features. *Mut Res* 531: 5-23.
- Carlson J, Locke AE, Flickinger M, Zawistowski M, Levy S, Myers RM, Boehnke M, Kang HM, Scott LJ, Li JZ
 et al. 2018. Extremely rare variants reveal patterns of germline mutation rate heterogeneity in
 humans. Nat Commun 9(1): 3753.
- 615 Cheng KC, Cahill DS, Kasai H, Nishimura S, Loeb LA. 1992. 8-Hydroxyguanine, an abundant form of 616 oxidative DNA damage, causes G>T and A>C substitutions. *J Biol Chem* **267**: 166-172.
- 617 Constantini D. 2014. Oxidative Stress and Hormesis in Evolutionary Ecology and Physiology. Springer,
 618 New York.
- Cook DE, Zdraljevic S, Roberts JP, Andersen EC. 2017. CeNDR, the *Caenorhabditis elegans* natural
 diversity resource. *Nucleic Acids Research* 45(D1): D650-D657.
- Cooke MS, Evans MD, Dizdaroglu M, Lunec J. 2003. Oxidative DNA damage: mechanisms, mutation, and
 disease. *FASEB J* 17: 1195-1214.
- 623 Cunningham RP. 1997. DNA repair: Caretakers of the genome? *Curr Biol* **7**: R576-R579.
- Denver DR, Dolan PC, Wilhelm LJ, Sung W, Lucas-Lledo JI, Howe DK, Lewis SC, Okamoto K, Thomas WK,
 Lynch M et al. 2009. A genome-wide view of *Caenorhabditis elegans* base-substitution mutation
 processes. *Proceedings of the National Academy of Sciences of the United States of America* 106(38): 16310-16314.
- Denver DR, Morris K, Kewalramani A, Harris KE, Chow A, Estes S, Lynch M, Thomas WK. 2004a.
 Abundance, distribution, and mutation rates of homopolymeric nucleotide runs in the genome
- 630 of Caenorhabditis elegans. Journal of Molecular Evolution **58**(5): 584-595.
- Denver DR, Morris K, Lynch M, Thomas WK. 2004b. High mutation rate and predominance of insertions
 in the *Caenorhabditis elegans* nuclear genome. *Nature* 430(7000): 679-682.
- 633 Denver DR, Morris K, Lynch M, Vassilieva LL, Thomas WK. 2000. High direct estimate of the mutation 634 rate in the mitochondrial genome of *Caenorhabditis elegans*. *Science* **289**(5488): 2342-2344.
- Denver DR, Wilhelm LJ, Howe DK, Gafner K, Dolan PC, Baer CF. 2012. Variation in base-substitution
 mutation in experimental and natural lineages of Caenorhabditis nematodes. *Genome Biology and Evolution* 4(4): 513-522.

Dollé MET, Snyder WK, Gossen JA, Lohman PHM, Vijg J. 2000. Distinct spectra of somatic mutations
 accumulated with age in mouse heart and small intestine. *Proc Nat Acad Sci USA* 97(15): 8403 8408.

- Drake JW, Charlesworth B, Charlesworth D, Crow JF. 1998. Rates of spontaneous mutation. *Genetics*148(4): 1667-1686.
- Eisen JA, Hanawalt PC. 1999. A phylogenomic study of DNA repair genes, proteins, and processes.
 Mutation Research 435(3): 171-213.
- Evans MD, Cooke MS. 2004. Factors contributing to the outcome of oxidative damage to nucleic acids.
 BioEssays 26: 533-542.
- Gao ZY, Moorjani P, Sasani TA, Pedersen BS, Quinlan AR, Jorde LB, Amster G, Przeworski M. 2019.
 Overlooked roles of DNA damage and maternal age in generating human germline mutations.
 Proceedings of the National Academy of Sciences of the United States of America 116(19): 9491 9500.
- Gray JM, Karow DS, Lu H, Chang AJ, Chang JS, Ellis RE, Marletta MA, Bargmann Cl. 2004. Oxygen
 sensation and social feeding mediated by a C-elegans guanylate cyclase homologue. *Nature* 430(6997): 317-322.
- 654 Guo C, McDowell IC, Nodzenski M, Scholtens DM, Allen AS, Lowe WL, Reddy TE. 2017. Transversions 655 have larger regulatory effects than transitions. *Bmc Genomics* **18**.
- Halldorsson BV, Palsson G, Stefansson OA, Jonsson H, Hardarson MT, Eggertsson HP, Gunnarsson B,
 Oddsson A, Halldorsson GH, Zink F et al. 2019. Characterizing mutagenic effects of
 recombination through a sequence-level genetic map. *Science* 363(6425): 364-373.
- Halligan DL, Keightley PD. 2009. Spontaneous mutation accumulation studies in evolutionary genetics.
 Annual Review of Ecology Evolution and Systematics 40: 151-172.
- Halliwell B, Gutteridge JMC. 2007. Free Radicals in Biology and Medicine. Oxford University Press,
 London.
- Hartman P, Ponder R, Lo HH, Ishii N. 2004. Mitochondrial oxidative stress can lead to nuclear
 hypermutability. *Mechanisms of Ageing and Development* 125(6): 417-420.
- Helbock HJ, Beckman KB, Shigenaga MK, Walter PB, Woodall AA, Yeo HC, Ames BN. 1998. DNA oxidation
 matters: The HPLC-electrochemical detection assay of 8-oxo-deoxyguanosine and 8-oxo guanine. Proc Nat Acad Sci USA 95: 288-293.
- Ishii N, Fujii M, Hartman PS, Tsuda M, Yasuda K, Senoo-Matsuda N, Yanase S, Ayusawa D, Suzuki K. 1998.
 A mutation in succinate dehydrogenase cytochrome *b* causes oxidative stress and ageing in
 nematodes. *Nature* 394: 694-697.
- Ishii N, Takahashi K, Tomita S, Keino T, Honda S, Yoshino K, Suzuki K. 1990. A methyl viologen sensitive
 mutant of the nematode *Caenorhabditis elegans*. *Mutation Research* 237(3-4): 165-171.
- Ishii T, Miyazawa M, Hartman PS, Ishii N. 2011. Mltochondrial superoxide anion (O2⁻⁻) inducible "*mev-1*"
 animal models for aging research. *BMB Reports* 44(5): 298-305.
- Ishii T, Miyazawa M, Onouchi H, Yasuda K, Hartman PS, Ishii N. 2013. Model animals for the study of
 oxidative stress from complex II. *Biochim Biophys Acta* 1827: 588-597.
- Ishii T, Yasuda K, Akatsuka A, Hino O, Hartman PS, Ishii N. 2005. A mutation in the *SDHC* gene of Complex
 Il increases oxidative stress, resulting in apoptosis and tumorigenesis. *Cancer Res* 65: 203-209.
- Ishii T, Yasuda K, Miyazawa M, Mitsushita J, Johnson TE, Hartman PS, Ishii N. 2016. Infertility and
 recurrent miscarriage with complex II deficiency-dependent mitochondrial oxidative stress in
 animal models. *Mech Ageing Dev* 155: 22-35.
- Joyner-Matos J, Bean LC, Richardson HL, Sammeli T, Baer CF. 2011. No evidence of elevated germline
 mutation acccumulation under oxidative stress in *Caenorhabditis elegans*. *Genetics* 189(4):
 1439-1447.

685 Katju V, Bergthorsson U. 2019. Old Trade, New Tricks: Insights into the Spontaneous Mutation Process 686 from the Partnering of Classical Mutation Accumulation Experiments with High-Throughput 687 Genomic Approaches. *Genome Biology and Evolution* **11**(1): 136-165. 688 Keightley PD, Caballero A. 1997. Genomic mutation rates for lifetime reproductive output and lifespan in Caenorhabditis elegans. Proceedings of the National Academy of Sciences of the United States of 689 690 America 94(8): 3823-3827. 691 Kessler MD, Loesch DP, Perry JA, Heard-Costa NL, Taliun D, Cade BE, Wang H, Daya M, Ziniti J, Datta S et 692 al. 2020. De novo mutations across 1,465 diverse genomes reveal mutational insights and 693 reductions in the Amish founder population. Proceedings of the National Academy of Sciences 694 **117**(5): 2560-2569. 695 Kim S-Y, Velando A. 2020. Attractive male sticklebacks carry more oxidative DNA damage in the soma 696 and germline. J Evol Biol 33: 121-161. 697 Konrad A, Brady MJ, Bergthorsson U, Katju V. 2019. Mutational landscape of spontaneous base 698 substitutions and small indels in experimental Caenorhabditis elegans populations of differing 699 size. Genetics 212(3): 837-854. 700 Konrad A, Thompson O, Waterston RH, Moerman DG, Keightley PD, Bergthorsson U, Katju V. 2017. 701 Mitochondrial Mutation Rate, Spectrum and Heteroplasmy in Caenorhabditis elegans 702 Spontaneous Mutation Accumulation Lines of Differing Population Size. Molecular Biology and 703 Evolution 34(6): 1319-1334. 704 Krašovec R, Richards H, Gifford DR, Hatcher C, Faulkner KJ, Belavkin RV, Channon A, Aston E, McBain AJ, 705 Knight CG. 2017. Spontaneous mutation rate is a plastic trait associated with population density 706 across domains of life. *PLoS Biology* **15**(8): e2002731. 707 Kreuz S, Fischle W. 2016. Oxidative stress signaling to chromatin in health and disease. *Epigenomics* 8(6): 708 842-862. 709 Liu HX, Zhang JZ. 2019. Yeast spontaneous mutation rate and spectrum vary with environment. Current 710 Biology 29(10): 1584-1591. 711 Lynch M, Hill WG. 1986. Phenotypic evolution by neutral mutation. *Evolution* **40**(5): 915-935. 712 Martin AP, Palumbi SR. 1993. Body size, metabolic rate, generation time, and the molecular clock. 713 Proceedings of the National Academy of Sciences of the United States of America 90(9): 4087-714 4091. 715 Matsuba C, Ostrow DG, Salomon MP, Tolani A, Baer CF. 2013. Temperature, stress and spontaneous 716 mutation in Caenorhabditis briggsae and Caenorhabditis elegans. Biology Letters 9(1). 717 McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D, Gabriel 718 S, Daly M et al. 2010. The Genome Analysis Toolkit: A MapReduce framework for analyzing next-719 generation DNA sequencing data. Genome Research 20(9): 1297-1303. 720 Messer PW. 2009. Measuring the Rates of Spontaneous Mutation From Deep and Large-Scale 721 Polymorphism Data. Genetics 182(4): 1219-1232. 722 Mukai T. 1964. Genetic structure of natural populations of Drosophila melanogaster. 1. Spontaneous 723 mutation rate of polygenes controlling viability. Genetics 50(1): 1-19. 724 Nei M, Li WH. 1979. Mathematical model for studying genetic variation in terms of restriction 725 endonucleases. Proceedings of the National Academy of Sciences of the United States of America 726 **76**(10): 5269-5273. 727 Paul C, Nagano M, Robaire B. 2011. Aging results in differential regulation of DNA repair pathways in 728 pachytene spermatocytes in the Brown Norway Rat. Biol Repro 85: 1269-1278. 729 Poetsch AR, Boulton SJ, Luscombe NM. 2018. Genomic landscape of oxidative DNA damage and repair 730 reveals regioselective protection from mutagenesis. Genome Biol 19: 215.

Richter C, Park JW, Ames BN. 1988. Normal oxidative damage to mitochondrial and nuclear DNA is
 extensive. Proceedings of the National Academy of Sciences of the United States of America
 85(17): 6465-6467.

Saxena AS, Salomon MP, Matsuba C, Yeh SD, Baer CF. 2019. Evolution of the mutational process under
 relaxed selection in *Caenorhabditis elegans*. *Molecular Biology and Evolution* 36(2): 239-251.

- Schrider DR, Houle D, Lynch M, Hahn MW. 2013. Rates and genomic consequences of spontaneous
 mutational events in *Drosophila melanogaster*. *Genetics* 194: 937-954.
- Senoo-Matsuda N, Yasuda K, Tsuda M, Ohkubo T, Yoshimura S, Nakazawa H, Hartman PS, Ishii N. 2001. A
 defect in the cytochrome b large subunit in complex II causes both superoxide anion
 overproduction and abnormal energy metabolism in *Caenorhabditis elegans. Journal of Biological Chemistry* 276(45): 41553-41558.
- 742 Sharp NP, Agrawal AF. 2012. Evidence for elevated mutation rates in low-quality genotypes. *Proceedings* 743 of the National Academy of Sciences of the United States of America **109**(16): 6142-6146.
- -. 2016. Low genetic quality alters key dimensions of the mutational spectrum. *PLoS Biology* **14**(3).
- 745Shigenaga MK, Gimeno CJ, Ames BN. 1989. Urinary 8-hydroxy-2'-deoxyguanosine as a biological marker746of *in vivo* oxidative DNA damage. *Proc Nat Acad Sci USA* **86**: 9697-9701.
- Sprague GF, Russell WA, Penny LH. 1960. Mutations affecting quantitative traits in the selfed progeny of
 doubled monoploid maize. *Genetics* 45(7): 855-866.
- 749Sterken MG, Snoek LB, Kammenga JE, Andersen EC. 2015. The laboratory domestication of750Caenorhabditis elegans. Trends in Genetics **31**(5): 224-231.
- Stoltzfus A. 2008. Evidence for a predominant role of oxidative damage in germline mutation in
 mammals. *Mut Res* 644: 71-73.
- Suzuki T, Kamiya H. 2017. Mutations induced by 8-hydroxyguanine (8-oxo-7,8-dihydroguanine), a
 representative oxidized base, in mammalian cells. *Genes and Environment* **39**: 2.
- Turrens JF. 2003. Mitochondrial formation of reactive oxygen species. *J Physiol* **552**: 335-344.
- Vassilieva LL, Lynch M. 1999. The rate of spontaneous mutation for life-history traits in *Caenorhabditis elegans. Genetics* 151(1): 119-129.
- Volkova NV, Meier B, Gonzalez-Huici V, Bertolini S, Gonzalez S, Vohringer H, Abascal F, Martincorena I,
 Campbell PJ, Gartner A et al. 2020. Mutational signatures are jointly shaped by DNA damage and
 repair. Nat Commun 11(1).
- Watterson GA. 1975. Number of segregating sites in genetic models without recombination. *Theoretical Population Biology* 7(2): 256-276.
- 763 Wright S. 1931. Evolution in Mendelian populations. *Genetics* **16**(2): 0097-0159.

764